

The Role of Quantum Optics in Next-Generation Secure Telecommunications

*¹Atsuwe, Bernard Aondofa, ²Atser, Akpenongun Roy, ³Gesa, Newton Felix and ⁴Ahemen, Iorkyaa



¹Department of Physics Education, Joseph Sarwuan Tarka University, Makurdi, Benue State, Nigeria.

²Department of Electrical/Electronics, Joseph Sarwuan Tarka University, Makurdi, Benue State, Nigeria.

³Department of Industrial Physics, Joseph Sarwuan Tarka University, Makurdi, Benue State, Nigeria.

⁴Department of Physics, Joseph Sarwuan Tarka University, P.M.B 2373, Makurdi, Benue State, Nigeria.

*Corresponding Author's Email: atsuwe.bernard@uam.edu.ng

ABSTRACT

Quantum optics, the study of light and its interactions with matter at the quantum level, has revolutionized modern telecommunications. By leveraging quantum phenomena such as entanglement, superposition, and photon manipulation, quantum optics enables ultra-secure communication, high-speed data transfer, and novel computing paradigms. The integration of quantum technologies into emerging communication infrastructures, particularly 6G networks, is becoming increasingly critical for overcoming the inherent limitations of classical communication and computation. In summary, quantum optics and quantum information have profoundly influenced modern physics and technology, where the field continues to bridge fundamental research and practical applications, promising further innovations in the future. The exponential growth of data traffic necessitates advancements in secure and high-speed communication technologies. Classical telecommunications rely on electromagnetic waves, but quantum optics introduces fundamentally new capabilities by exploiting quantum states of light. This paper explores the fundamental principles of quantum optics, its applications in quantum key distribution (QKD), quantum teleportation, and quantum repeaters, and its impact on next-generation telecommunications. We also discussed challenges and future prospects of the quantum optics in telecommunications.

Keywords:

Quantum optics,
Quantum communication,
QKD,
Entanglement,
Quantum teleportation,
Optical fibers.

INTRODUCTION

Quantum optics has evolved from a specialized area within atomic, molecular, and optical physics into a vibrant and interdisciplinary field that bridges multiple branches of physics. Its growing prominence is largely due to its central role in quantum information science and its connections to ultra-cold quantum gases, which have revealed fascinating links to condensed-matter physics. Despite these advancements, quantum optics retains a distinctive foundation of core principles, which this concise paper aims to elucidate. At its heart, quantum optics explores the most elementary quantum systems, such as light and atoms with few energy levels. These systems exemplify quantum mechanics in its purest form, free from the complexities of larger-scale phenomena. Remarkably, quantum optics often brings to life the theoretical experiments envisioned by the pioneers of quantum mechanics. For the foreseeable future, the field will remain a cornerstone of quantum physics, centered

on elegantly simple yet profound ideas and experiments (Ulf, 2010).

A defining feature of quantum optics is the seamless interplay between theory and experimentation. While most literature focuses on theoretical concepts, many of these ideas are rooted in experimental observations or have, in turn, driven innovative experiments. Another strength of the field lies in its accessibility to individual researchers or small teams. A single scientist or a modest group can design and conduct an entire experiment, while a theorist can tackle significant problems with minimal tools—sometimes just pencil and paper. As John A. Wheeler aptly noted in Ulf, 2010 that "It is gratifying to know that computers can solve the problem, but true understanding comes from human insight."

The year 1900 marked a pivotal moment in physics when Max Planck proposed that the energy of the light field is quantized. This groundbreaking idea laid the foundation for quantum mechanics, which has since become a

cornerstone of modern physics. The invention of the laser in 1960 further revolutionized the field, providing a precise tool for generating coherent monochromatic light. This advancement propelled quantum optics forward, driven by a synergy of experimental and theoretical research. The contributions of Sir Peter Knight, a trailblazer in quantum optics, known for his pioneering work, influential reviews, and textbooks (Knight, & Allen, 1983; Gerry, & Knight, 2005) is regarded as being key to quantum optics. Quantum optics has played a critical role in the advancement of quantum technology and quantum information, with Knight being one of its key architects. The field has enabled precise tests of quantum mechanics, such as the observation of "quantum jumps," which highlight the discontinuous nature of quantum measurements (Cook, & Kimble, 1985; Knight, Loudon, & Pegg, 1986; Bergquist, Hulet, Itano, & Wineland, 1986). Among its central concepts is the coherent state, introduced by Schrödinger and later expanded by Glauber and Sudarshan to analyze high-order light coherences (Glauber, 1963; Sudarshan, 1963). Squeezed light, which redistributes quantum noise between quadratures, has also been instrumental, finding applications in precision measurements like gravitational wave detection and noiseless communication (Loudon, & Knight, 1987; Asasi, et al., 2013; Yuen, & Shapiro, 1978). The Jaynes-Cummings model (JCM) (Jaynes, & Cummings, 1963; Shore, & Knight, 1963), describing the interaction between a two-level atom and a single electromagnetic field mode, has been a powerful framework for exploring nonclassical phenomena. For instance, the quantization of the field leads to collapses and revivals of Rabi oscillations, demonstrated in cavity quantum electrodynamics (QED) experiments (Eberly, Narozhny, & Sanchez-Mondragon, 1980; Rempe, Walther, & Klein, 1987; Meekhof, Monroe, King, Itano & Wineland, 1996). Information-theoretic studies of the JCM revealed that a cavity field initially in a coherent state evolves into a superposition state at specific interaction times (Plenio, & Knight, 1998; Phoenix, & Knight, 1988; Gea-Banacloche, 1990; Lv et al., 2017). The exponential growth in the complexity of quantum systems poses challenges for exact calculations, prompting Richard Feynman to propose simulating such systems using controllable quantum entities (Feynman, 1982). Superposition in qubits leads to entanglement, enabling nonclassical information processing. Quantum computing, for example, leverages these principles to solve problems like integer factorization efficiently, as demonstrated by Shor's algorithm (Shor, 1997). Quantum mechanics also enhances secure communication; Bennett and Brassard's 1984 protocol (Bennett & Brassard, 1984) used superposition states of light to create a one-time pad, a highly secure encryption method.

A major challenge in quantum technology is decoherence, caused by unwanted environmental

interactions. Research into Markovian and non-Markovian open quantum systems has explored decoherence mechanisms. Knight and Plenio investigated quantum decoherence using quantum jump methods (Plenio, & Knight, 1998) and developed strategies to mitigate it, such as phase control (Paspalakis, & Knight, 1998) and decoherence-free subspaces (Beige, et al., 2000). Quantum error correction, requiring extensive entangled resources (Shor, 1995), and fault-tolerant quantum computation (Steane, 1999) are also vital for overcoming decoherence. Entanglement, a hallmark of quantum mechanics, describes correlations where local systems may exhibit more uncertainty than the global system. This phenomenon, linked to the Einstein-Podolsky-Rosen (EPR) paradox (Einstein, Podolsky, & Rosen, 1935), has practical applications. For instance, Ekert's protocol (Ekert, 1991) uses entangled states for secure key distribution. Quantum teleportation, another application, relies on entanglement to transmit quantum states, with entanglement distillation protocols (Bennett, et al., 1996; Bose, Vedral, & Knight, 1999) and swapping (Zukowski, et al., 1993; Bose, Vedral, & Knight, 1998) enhancing its feasibility. Knight and colleagues advanced the quantification of entanglement (Ekert, & Knight, 1995; Vedral, et al., 1997; Vedral, & Plenio, 1998; Lee, et al., 2000), contributing to its theoretical understanding. Atom optics, another branch of quantum optics, exploits quantum coherence in atomic motion. Techniques like Ramsey interferometry, based on atomic internal state coherences, have led to atomic clocks and precision sensors (Deutsch, 1985).

Quantum optics and quantum information have profoundly influenced modern physics and technology, with Sir Peter Knight's work being central to these developments. The field continues to bridge fundamental research and practical applications, promising further innovations in the future. The exponential growth of data traffic necessitates advancements in secure and high-speed communication technologies. Classical telecommunications rely on electromagnetic waves, but quantum optics introduces fundamentally new capabilities by exploiting quantum states of light. This paper examines how quantum optics enhances telecommunications through: Quantum Key Distribution (QKD) for unbreakable encryption, Quantum Teleportation for state transfer without physical transmission, and Quantum Repeaters for long-distance entanglement distribution. Quantum optics, a field that explores the interaction between light and matter at the quantum level, has revolutionized the field of telecommunications. By harnessing the principles of quantum mechanics, quantum optics enables the development of advanced technologies that can transform the way we communicate. Quantum optics is based on the principles of quantum mechanics, which

describe the behavior of particles at the atomic and subatomic level. In the context of telecommunications, quantum optics focuses on the manipulation of light, specifically photons, to transmit information securely and efficiently.

The rapid evolution of global telecommunications demands security and efficiency beyond the capabilities of classical systems. While classical cryptography and signal processing have advanced considerably, they remain vulnerable to computational breakthroughs and eavesdropping. Quantum optics, rooted in the quantum mechanical behavior of light, offers a paradigm shift by exploiting properties such as entanglement and superposition to achieve theoretically unhackable communication and novel networking architectures. Existing reviews often summarize quantum optical principles and applications without critically addressing the gap between theoretical promise and practical implementation. This paper moves beyond mere summarization to synthesize insights from the literature, focusing on the challenges that hinder real-world

deployment and the solutions proposed to overcome them. We argue that the future of quantum telecommunications depends not only on advancing core technologies like QKD and quantum repeaters but also on solving integration, scalability, and noise-related issues.

Theoretical Foundations of Quantum Optics

Some key concepts in quantum optics according to (Gisin, et al., 2002; Bell, 1964) include Quantization of Light which is the idea that light is composed of discrete packets, or photons, rather than being a continuous wave, Wave-Particle Duality that entails the ability of light to exhibit both wave-like and particle-like behavior, Superposition which is the ability of quantum systems to exist in multiple states simultaneously and Entanglement that is the phenomenon where two or more quantum systems become correlated in such a way that the state of one system cannot be described independently of the others.

Table 1: Key Concepts in Quantum Optics (Field, 2025)

S/N	Concept	Description
1.	Quantization of Light	Light is composed of discrete packets, or photons
2.	Wave-Particle Duality	Light exhibits both wave-like and particle-like behavior
3.	Superposition	Quantum systems can exist in multiple states simultaneously
4.	Entanglement	Quantum systems become correlated in such a way that the state of one system cannot be described independently of the others

Quantum Nature of Light

Light exhibits both wave-like and particle-like (photon) behavior. Photons, the quanta of light, are fundamental to quantum optical processes.

Some key quantum optical concepts include (Shor, 1997):

- i. Photon Statistics: Described by Bose-Einstein distribution.
- ii. Coherent States: Laser light approximates coherent states.
- iii. Squeezed Light: Reduced noise in one quadrature at the expense of another.

Initially, the quantum behavior of light can appear peculiar and confusing. The duality of light existing as distinct packets of energy, massless particles that carry momentum, and fundamental entities exhibiting wave-like properties can feel as fantastical as a narrative from "Alice in Wonderland". For numerous individuals, delving into this physics discipline is as mesmerizing as Lewis Carroll's renowned book. Reflecting on the contributions of iconic figures and esteemed scientists like Einstein, Planck, and Compton, examining light's quantum character reveals a compelling story of how intelligent interpretation of minor observations sparked the most crucial scientific advancements of the past 150 years. From the electronics transformation of the

twentieth century to upcoming innovations in solar power and space research, the quantum aspect of light promises a domain of significant outcomes, housing some of the most captivating realities of our era (Bouchee et al., 2023).

Quantum Superposition

A quantum system can exist in multiple states simultaneously until measured. In optics, this allows photons to be in superposition states, enabling quantum computing and communication protocols.

The concept of superposition in its initial quantum physics form relates to waves, stemming from the superposition principle established in mathematics for solutions to a linear differential equation. Within quantum mechanics, the mathematical framework for quantum physics, abstract quantum states and their superpositions are not always solutions of such an equation. As a generalization of the principle proven for linear differential equations, the quantum-mechanical superposition principle is accepted as a fundamental postulate. It is not derivable from more basic concepts and therefore requires no proof. As per this principle, if a quantum system can exist in one of two states, it can also exist in a linear combination of these states. This

combined state is also known as a quantum superposition. (Guang-Liang, 2023)

Consequently, in general terms, quantum superpositions can be classified into two types: the superposition of waves, which originates from the mathematical superposition principle for solutions of a linear differential equation, and the superposition of abstract quantum states. These quantum states can be linked to either a single quantum entity or a system of multiple quantum objects.

Quantum Entanglement

When two or more photons become entangled, their quantum states are intrinsically linked, regardless of distance. This phenomenon is crucial for quantum teleportation and secure communication. Entangled photons exhibit correlated properties regardless of distance, forming the basis for quantum communication. The appearance of the combined term "quantum entanglement" in published literature seems to trace back to the mid-to-late 1980s (Ghirardi, et al., 1987). This occurred several years after pioneering optical experiments on quantum entanglement were conducted by Aspect and colleagues in the early 1980s (Aspect, et al., 1981, 1982a, 1982b). Currently, the domain of quantum entanglement is vast and is approximately separated into three primary subfields as: quantum cryptography, quantum teleportation, and quantum computing. Quantum communications falls under quantum cryptography. On paper, based on citation analysis, these subfields have been strongly shaped by ideas and concepts originating from the philosophical approach to quantum entanglement (Duarte, 2022). Conversely, also from a citation perspective, recognition of the physics-based path has been exceptionally limited (Duarte, 2022). This near-total neglect has continued

despite the critically important probability amplitude for quantum entanglement, which was identified devoid of philosophical debate: being reintroduced into the mainstream quantum entanglement literature by Greenberger et al. in 1990.

The evolution of quantum physics theory is highly complex, as it encompasses the work of many different physicists. Therefore, in addition to clarifying why faster-than-light communication is unachievable, Yihe-Xue, 2021 moved further to propose theories on superposition mechanisms. This he did because any deeper investigation into contemporary quantum entanglement theory demands a strong foundational knowledge. A structured historical account can efficiently conserve time and lower the probability of errors. For humanity to advance further, deciphering the enigmas in quantum physics is essential. Although quantum entanglement cannot be utilized for faster-than-light communication, its impact on other domains, such as information encryption, remains highly significant (Yihe-Xue, 2021).

Quantum entanglement is a unique occurrence that exists solely within quantum physics. No analogous phenomenon exists in the annals of classical physics. The concept was initially introduced by Erwin Schrödinger following the Einstein-Podolsky-Rosen paradox (Manik, et al., 2019). Motivated by the thought that experiments in the quantum Mechanical description of physical reality can be considered complete (Albert, et al., 1935), observed that after particles interact, they can only be characterized as a unified whole, not as independent entities; they become entangled. Spin is a particle property that exhibits quantum entanglement. Therefore, measuring spin is crucial for advanced quantum research (Yihe-Xue, 2021).




Figure 1: Quantum Entanglement (Manik, et al., 2019)

Beam Splitters and Interferometers

The beam splitter has served multiple functions across various optics disciplines. For instance, in quantum information science, it is crucial for teleportation, Bell

state measurements, entanglement generation, and fundamental photon research. This is critical for manipulating quantum states in optical circuits (Fig. 2).




Figure 2: (a) Schematic Diagram of the Mach-Zehnder Interferometer with a Quantum Beam Splitter BS_2 . (b) The Quantum Circuit that Describes the Evolution of the Ancilla and the Photon in the Interferometer (Filgueiras, et al., 2012)

No-Cloning Theorem

Quantum information cannot be perfectly copied, ensuring security in quantum communication by preventing eavesdropping. These principles form the foundation for advanced quantum communication technologies.

Quantum Optical Technologies in Telecommunications

Quantum Key Distribution (QKD)

Quantum Key Distribution (QKD) leverages quantum optics to enable ultra-secure communication. The most well-known protocol, BB84, uses polarized photons to generate encryption keys.

Quantum key distribution (QKD) is a component of quantum cryptography. It refers to a collection of protocols that enable the expansion of an initial secret key, shared solely by the two communicating parties, into a larger key. Given reasonable assumptions, the information leakage to external parties from the newly generated bits can be constrained as narrowly as needed, thereby offering a method for secure symmetric key generation. Apart from correct protocol execution, the assumptions are confined to the implementation's physics: (i) the participants' equipment does not leak information externally (i.e., a security perimeter exists), a standard prerequisite for any cryptosystem; (ii) both participants can access a genuine randomness source, such as a properly implemented quantum random number generator (QRNG); and (iii) the known laws of quantum mechanics are universally applicable, thus limiting any potential adversaries (Scarani, 2009). In this regard, QKD security is grounded in physical laws. QKD

protocols are therefore provably absolutely secure in a mathematical sense, a point we will elaborate on later. No imaginable attack, whether classical or quantum, could compromise the system, regardless of the resources employed. This stands in contrast to other modern key distribution protocols that rely on mathematically unverified assumptions for their security. A key example is the widely used Diffie-Hellman protocol, which depends on the computational difficulty of the discrete logarithm problem for security. If the computational complexity is actually lower than presumed, the algorithm would be rendered ineffective, and any past Diffie-Hellman exchange could be decoded, including historical ones. Indeed, the standard computational complexity assumption for this problem fails in the quantum realm. A quantum computer could break Diffie-Hellman and other prevalent protocols like RSA and elliptic curve cryptography using Shor's algorithm (Shor, 1994; Shor, 1997). Algorithms that can be executed more efficiently on a quantum computer than on classical ones form the subject of quantum computing. Classical algorithms designed to resist Shor's algorithm, or quantum computing in general, are the focus of post-quantum cryptography.

QKD protocols necessitate the capability to generate, control, transmit, and detect signals at the quantum level. This constitutes a highly challenging technology, though the transition to real-world application was relatively swift. It was first demonstrated in a laboratory as a proof of concept in 1989 (Bennett, & Brassard, 1989) and became a commercial product as early as 2002 (ID Quantique, ND). Since then, it has been deployed in practical scenarios numerous times, including within

classical communication networks, and several firms market QKD devices in the security industry. The principal distinctions between QKD and equivalent algorithms (i.e., symmetric key) in traditional cryptography are that it addresses the secret key agreement problem in an unconditionally secure manner (under the aforementioned assumptions). QKD accomplishes this, however, by requiring a quantum channel—a communication link capable of transmitting quantum correlations—that must be physically established between the two parties (Vincente, Jesus, & Momtchil, 2016). It also needs an authenticated classical channel with integrity protection, but this is readily available in modern communication infrastructures. In comparison, the conventional method of secret key distribution only requires an algorithm (e.g., Diffie-Hellman) operating on a computer and an accessible classical communication channel. Cryptographically, a pair of QKD systems can be viewed as a seeded, distributed, and correlated randomness source, with the additional assurance that only a limited, controllable

amount of information is leaked outside the security boundaries of the two legitimate parties. Another perspective is to see it as an extension of the security perimeter of the two parties' devices to include the connecting communication line. QKD Works in such a way that a sender (Alice) transmits photons in random polarization states, a receiver (Bob) measures the photons using randomly chosen bases and eavesdropping (by Eve) disturbs the quantum states, alerting Alice and Bob to potential interception (Vincente, Jesus, & Momtchil, 2016).

The advantages of QKD amongst others are: Unbreakable Security Based on quantum principles, QKD detects any intrusion, Future-Proof Resistant to attacks from quantum computers e.t.c. Real-world implementations include China's Micius satellite and commercial QKD networks by companies like ID Quantique and Toshiba. QKD enables secure communication using quantum principles: BB84 Protocol: Uses polarization states for key exchange. E91 Protocol: Relies on entangled photons.




Figure 3: Framework of the Quantum key Distribution (QKD) Network (Tsai,2021)

Quantum Key Distribution (QKD) principles, their accompanying challenges and solutions are summarized as:

- i. Principle: Uses quantum states to detect eavesdropping.
- ii. Challenges: Limited transmission distance, high cost, integration with existing infrastructure.
- iii. Solutions: Satellite-based QKD (e.g., Micius), hybrid classical-quantum networks, improved photon sources and detectors.

Table 2: Comparison of QKD Protocols (Field, 2025)

S/N	Protocol	Basis Used	Security Mechanism
1.	BB84	Polarization	Heisenberg Uncertainty
2.	E91	Entanglement	Bell's Theorem

Quantum Teleportation

For many, quantum physics is among the most captivating yet baffling areas of modern physics. Despite its counter-intuitive nature, quantum physics is anticipated to greatly influence the technologies underpinning our future communication and online security (Wehner, Elkouss & Hanson, 2018; Wei S-H *et al* 2022; Sridhar, Ashwini & Tabassum, 2023). From the many quantum physics ideas, quantum teleportation is one of the most compelling. This procedure is vastly different from matter teleportation, a notion popularized by various fictional gadgets like the “Star Trek” transporter (Krauss, 1995). Matter teleportation involves moving people or objects between locations, which is not feasible with our present comprehension of physical laws. (Fitzgerald, Emonts, & Tura, 2024).

Quantum teleportation, however, is achievable, and unlike fictional matter teleportation where atoms and molecules are relocated, it is used to transfer quantum information between two sites over distances of 1000 km or more (Yin *et al.*, 2017). In the future, quantum teleportation will be instrumental in creating quantum networks, a prerequisite for more complex algorithms like quantum key distribution, a core protocol for quantum-secure communication (Bennett, & Brassard, 2014; Zapatero *et al.*, 2023). Hence, it is vital that the basics, potential, and constraints of quantum teleportation are communicated clearly and innovatively to the upcoming generation of quantum physicists in schools (Singh, Levy, & Levy, 2022) and that instructors receive adequate training to support this (Sutirini *et al.*, 2023). The second quantum revolution, which foresees progress in communication (partly based on quantum teleportation), sensors, computers, and simulations,

"poses new challenges for the new emerging workforce who will develop these quantum technologies or work with them" (Greinert *et al.*, 2023). However, tackling the technical obstacles is only one aspect. There are many challenges in educating students initially, such as securing qualified teaching staff and addressing insufficient financial investment in essential educational infrastructure and equipment (Aiello *et al.*, 2021). Moreover, there is a pressing need for this education as quantum physics increasingly permeates daily life, especially given the prevalence of pseudo-knowledge and misinformation (Introne *et al.*, 2018), which has also impacted quantum science (Meyer *et al.*, 2023). Quantum teleportation allows the transfer of quantum states between distant particles using entanglement.

Process of Quantum Teleportation

- i. Entanglement Distribution: Two entangled photons are shared between sender (Alice) and receiver (Bob).
- ii. Bell-State Measurement: Alice performs a joint measurement on her photon and the quantum state to be teleported.
- iii. Classical Communication: Alice sends measurement results to Bob via a classical channel.
- iv. 4. State Reconstruction: Bob applies corrections to his photon to replicate the original quantum state.

Applications in Telecommunications

Enables secure quantum networks and Facilitates long-distance quantum communication. Quantum teleportation transfers quantum states between distant nodes using entanglement and classical communication (Figure 2). (Fitzgerald, Emonts & Tura, 2024)




Figure 4: A schematic Demonstration of the Quantum Teleportation Protocol

Figure 4 shows a schematic demonstration of the quantum teleportation protocol. The circles represent particles, the solid line represents the quantum communication channel (entanglement), and the dashed line represents the classical communication channel. In the (a) above, Alice and Bob Initially share some

entanglement via particles 1 and 2. Alice has a third particle in which a quantum state ψ is encoded. While in (b) Alice subjects particles 2 and 3 to a Bell-state measurement (BSM), which generates two bit of classical information j , that distinguish between the four possible measurement outcomes. Alice communicates this

information to Bob using a classical channel, whereupon Bob uses his knowledge of j to apply the appropriate unitary operator U_j and ensure his particle now encodes the state \square (Travis, 2014).

The principles, challenges and solutions can be summarized as:

- i. Principle: Transfers quantum states using entanglement and classical communication.
- ii. Challenges: Fidelity loss, decoherence, need for quantum memory.
- iii. Solutions: Entanglement purification advanced Bell-state measurements, error correction codes.

Quantum Repeaters

Quantum repeaters are instruments that increase the distance of quantum channels. They do this by disseminating entangled states among far-apart nodes of a quantum network. Entanglement acts as a resource for long-range quantum communication. A comprehensive quantum network using quantum repeater technology is built by combining and linking numerous short-range quantum communication segments. These individual short-range links are each precisely adjusted to be shorter than the attenuation length of the quantum communication channel (Das, Rahman, & Majumdar, 2021). The quantum repeaters are situated as intermediate units between nodes of neighboring links, effectively connecting two communication segments. The quantum repeater executes entanglement swapping between nodes of adjacent links, thus generating entanglement between the endpoints of these links. This procedure is repeated across the whole transmission path, finally resulting in the entanglement of the sender and receiver nodes (Ruihong, & Ying, 2019; Gisin, & Thew, 2007). As a result, a strong quantum communication link is formed between them, enabling direct, secure, and effective quantum communication. This capacity to prolong quantum entanglement over large distances is crucial for the real-world application of quantum communication and quantum key distribution globally.

The key elements of a quantum repeater include: quantum entanglement switching for exchanging entangled states between nearby nodes, quantum memory for storing quantum states for efficient retrieval when needed, and quantum entanglement purification for improving the quality of the entangled states. These elements face some constraints due to flaws in the source of entangled particles, the quantum operations, and the connecting communication channels (Herbst et al. 2015). Notwithstanding these constraints, quantum repeaters have been shown to work in experiments, like those by Herbst et al., 2015 that employed a quantum repeater to teleport an entangled state, a photon, between La Palma and Tenerife in the Canary Islands, a distance of roughly 143 km (Herbst et al., 2015). The entanglement swapping experiment utilized two

polarization-entangled photon pairs produced in two separate spontaneous parametric down-conversion (SPDC) sources employing a non-linear crystal, “ β ”-barium borate (BBO) (Herbst et al. 2015).

The quantum entanglement purification protocol is a vital element in quantum repeaters and is needed for first-generation or near-future quantum repeaters (Muralidharan et al., 2016). It is the method of refining high-quality entanglement from lower-quality entanglement. Its purpose is to guarantee that entangled states retain high fidelity during transmission (Bennett et al., 1996), thereby offsetting the fidelity loss caused by noise or flaws in the communication channel. The two purification methods used in this study are: Bennett's protocol (Bennett et al., 1996) and Deutsch's protocol (Deutsch et al., 1996). Performance factors evaluated for the purification protocols include the fidelity of the purified Bell pair and the success probability (Krastanov, Albert, & Jiang, 2019). All protocols aim to produce shorter circuits that attain higher success rates and better final fidelity (Krastanov, Albert, & Jiang, 2019). Quantum repeaters are essential for future quantum communication technologies like the quantum internet (Briegel et al. 1998; Gisin, & Thew, 2007). They will increase the range of transmission links to a global, inter-continental scale, driving the future of a worldwide quantum network. A significant effort in this area involves attempting to integrate quantum repeaters into current optical fibre infrastructure (Rabbie et al., 2022). Experiments, such as those by Hu et al., 2021 where they effectively disseminated a pair of polarization-entangled states over a notable 11 km distance using noisy multicore fibre cables (Hu et al. 2021), build assurance in the viability of setting up quantum communication links within existing optical fibre systems. However, realizing practical, full-scale quantum repeaters in the real world remains a major technological hurdle. Considerable ongoing research is examining both the individual parts and the complete architecture of quantum repeater systems.

This research details the full-scale architecture of a quantum repeater, designed and realized using quantum circuits run on a quantum computer. We employed this quantum repeater implementation to investigate quantum entanglement purification (Mugambi, & Okeng'o, 2023). A major challenge in quantum communication is photonic loss over long distances. Classical optical repeaters amplify signals, but quantum signals cannot be copied due to the no-cloning theorem. How Quantum Repeaters Work: Use entanglement swapping to extend entanglement over long distances. Employ quantum memory to store and retrieve quantum states. Benefits of quantum repeaters in telecommunications include, Extending the range of quantum networks beyond direct fiber limits (~100 km) and it is essential for a global quantum internet amongst others.

Summarily, quantum repeaters can be presented in the form as:

- i. Principle: Extend entanglement range via swapping and purification.
- ii. Challenges: Photon loss in fibers, slow entanglement distribution, quantum memory coherence time.

- iii. Solutions: Multi-node networks, atomic ensemble memories, all-photonic repeaters.

Critical Challenges and Synthesis of Solutions

The following table summarizes major challenges and solution pathways derived from the literature:

Table 3: Major Challenges and Solution Pathways Derived from Literature

S/N	Challenge Category	Specific Issues	Proposed Solutions	Research Gaps
1.	Technical Limitations	Photon loss, decoherence, low detector efficiency	Squeezed light, improved single-photon detectors, fault-tolerant designs	Long-lived quantum memory, high-rate entanglement sources
2.	Scalability & Cost	Expensive hardware, limited range, network integration	Modular quantum repeaters, chip-based photonics, shared infrastructure	Cost-effective QKD systems, scalable quantum network protocols
3.	System Integration	Compatibility with classical networks, standardization	Hybrid networks, software-defined quantum networking	Interoperability standards, quantum-classical interface design
4.	Security Assurance	Side-channel attacks, implementation flaws	Device-independent QKD, security certification frameworks	Real-world attack resilience, quantum firewall development

Impact of Quantum Optics on Next-Generation Telecommunications

Quantum optics is revolutionizing telecommunications in several ways:

- i. Ultra-Secure Communication: QKD ensures military-grade security for governments, banks, and enterprises.
- ii. High-Speed Quantum Networks: Quantum entanglement enables faster-than-classical information transfer in distributed systems.

- iii. Quantum Internet: A future global quantum internet could connect quantum computers and sensors with unprecedented efficiency.

- iv. Post-Quantum Cryptography: Quantum-resistant encryption methods are being developed to counter future quantum hacking threats.

Photons degrade in optical fibers; quantum repeaters extend entanglement distribution via entanglement swapping (Fig. 4).




Figure 5: A schematic Diagram of the Entanglement Swapping Process

Role of Quantum Optics in Advanced Telecommunications

Quantum optics plays a crucial role in advanced telecommunications by enabling (Gisin et al., 2002; Krijfenburg-lewerissa, Brinkman, & Joolingen, 2017):

- i. Secure Communication: Quantum cryptography, also known as quantum key distribution (QKD), uses quantum optics to create secure communication channels. QKD relies on the

principles of quantum mechanics to encode and decode messages, making it virtually un-hackable.

- ii. Quantum Telecommunication Networks: Quantum optics enables the development of quantum telecommunication networks, which can transmit quantum information over long distances. These networks have the potential to revolutionize the way we communicate, enabling secure and efficient communication over long distances.

Table 4: Comparison of Quantum Optics-Based Systems and Classical Systems

S/N	Feature	Quantum Optics-based Systems	Classical Systems
1.	Security	Enhanced security through quantum cryptography	Vulnerable to hacking and eavesdropping
2.	Efficiency	Potential for faster data transmission rates	Limited by classical physics
3.	Reliability	Potential for more reliable data transmission	Prone to errors and interference

Applications of Quantum Optics in Telecommunications

Quantum optics, a branch of physics that deals with the behavior of light and its interactions with matter at the quantum level, has numerous applications in telecommunications. It has a wide range of applications where by harnessing the principles of quantum mechanics, quantum optics can enable advanced telecommunications systems that are more secure, efficient, and reliable. As research and development in this field continue to advance, we can expect to see widespread adoption of quantum optics-based systems in telecommunications. Some of the key applications of quantum optics in telecommunications include (Nielsen, & Chuang, 2010):

- i. Quantum Key Distribution (QKD): One of the most significant applications of quantum optics in telecommunications is Quantum Key Distribution (QKD) (Gisin et al., 2002). QKD uses quantum optics to create secure communication channels, enabling secure data transmission over long distances. QKD systems rely on the principles of quantum mechanics to encode and decode messages, making them virtually un-hackable.
- ii. Quantum Cryptography: Quantum optics enables secure communication channels, making it ideal for sensitive data transmission (Nielsen, & Chuang, 2010). Quantum cryptography, which uses quantum optics to encode and decode messages, provides an additional layer of security for data transmission. This is particularly important for applications such as financial transactions, military communications, and sensitive data transfer.
- iii. Optical Quantum Memory: Optical quantum memory is a critical component of quantum telecommunication networks, enabling the storage and retrieval of quantum information (Bennett, & Brassard, 1984). This technology has the potential to enable the development of quantum repeaters, which can extend the range of quantum communication systems.
- iv. Quantum Computing: Quantum optics is used to develop quantum computers, which have the potential to solve complex problems that are beyond the capabilities of classical computers.
- v. Quantum Cryptography: Quantum cryptography uses quantum optics to encode and decode messages, making it virtually un-hackable. This technology has numerous applications in secure

communication, including secure data transmission over long distances.

- vi. Quantum Telecommunication Networks: Quantum optics enables the development of quantum telecommunication networks, which can transmit quantum information over long distances (Ladd et al., 2010). These networks have the potential to revolutionize the way we communicate, enabling secure and efficient communication over long distances.
- vii. Spectroscopy: Quantum optics is used in spectroscopy to study the properties of atoms and molecules. By understanding the principles of quantum optics, we can develop new technologies and applications that take advantage of the unique properties of light at the quantum level.

Benefits of Quantum Optics in Telecommunications

The benefits of quantum optics in telecommunications include:

- i. Enhanced Security: Quantum optics enables secure communication channels, making it ideal for sensitive data transmission.
- ii. Increased Efficiency: Quantum optics can enable faster data transmission rates, making it ideal for high-speed data transfer applications.
- iii. Improved Reliability: Quantum optics can enable more reliable data transmission, reducing errors and improving overall system performance.

Quantum Optics Experiments

Some notable quantum optics experiments include (Ladd et al., 2010):

- i. Double-Slit Experiment: A classic experiment that demonstrates the wave-particle duality of light.
- ii. Hong-Ou-Mandel Experiment: An experiment that demonstrates the phenomenon of quantum entanglement.
- iii. Quantum Eraser Experiment: An experiment that demonstrates the ability to retroactively change the outcome of a measurement.

Quantum Optics Technologies

Some notable quantum optics technologies include:

- i. Lasers: Lasers are a key technology in quantum optics, enabling the manipulation of light at the quantum level.
- ii. Optical Fibers: Optical fibers are used to transmit quantum information over long distances.

- iii. Photo detectors: Photo detectors are used to detect and measure the properties of photons.

Challenges and Future Directions

Challenges: While quantum optics has the potential to revolutionize telecommunications, there are several challenges that need to be addressed. These include:

- i. Scalability: Quantum optics-based systems need to be scalable to be practical for widespread adoption.
- ii. Cost: Quantum optics-based systems are currently expensive, making them less competitive with classical systems.
- iii. Interoperability: Quantum optics-based systems need to be interoperable with existing classical systems.
- iv. Photon Loss and Decoherence: Optical fibers attenuate quantum signals over long distances.
- v. Technological Limitations: High-precision quantum detectors and memories are still under development.
- vi. Noise and Error Correction: Quantum optics-based systems are prone to noise and errors, which need to be corrected.
- vii. Quantum Control: Maintaining control over quantum systems is essential for practical applications.

Future Prospects

- i. Satellite-Based QKD: Expanding secure communication via satellites (e.g., China's Micius).
- ii. Hybrid Quantum-Classical Networks: Integrating quantum and classical systems for practical deployment.
- iii. Advances in Quantum Repeaters: Overcoming distance limitations for global quantum networks.
- iv. Integrated Photonics: Scalable quantum chips.

Future Research Directions

Based on our synthesis, the following areas present significant opportunities for original research:

- i. Development of Long-Coherence Quantum Memories: Materials and designs that extend storage time beyond milliseconds are critical for practical quantum repeaters.
- ii. Integrated Quantum Photonics: On-chip quantum light sources, detectors, and circuits to reduce cost and improve scalability.
- iii. Quantum Network Protocols: New protocols for routing, error management, and network discovery in large-scale quantum internet.
- iv. Post-Quantum Hybrid Systems: Integration of QKD with post-quantum cryptographic algorithms for layered security.
- v. Standardization and Certification: Establishing global standards for quantum communication hardware and software.

CONCLUSION

Quantum optics is a rapidly advancing field that has revolutionized our understanding of light and its properties. From quantum computing to quantum cryptography, quantum optics has enabled the development of many new technologies and applications. As research and development in this field continue to advance, we can expect to see widespread adoption of quantum optics-based systems in a wide range of fields. Quantum optics is transforming telecommunications by enabling ultra-secure, high-speed, and long-distance quantum communication. From QKD and quantum teleportation to quantum repeaters, these technologies are paving the way for a quantum internet. Quantum optics has numerous applications in telecommunications, including secure communication, quantum telecommunication networks, and optical quantum memory. By harnessing the principles of quantum mechanics, quantum optics enables the development of advanced telecommunication systems that are more secure, efficient, and reliable. Despite challenges like photon loss and scalability, ongoing research and technological advancements promise a future where quantum communication becomes mainstream. As research and development in this field continue to advance, we can expect to see widespread adoption of quantum optics-based systems in telecommunications and as we move toward next-generation networks, quantum optics will play a pivotal role in shaping the future of global telecommunications.

REFERENCES

- Aiello, C. D. *et al* (2021) Achieving a quantum smart workforce *Quantum Sci. Technol.* **6** 030501
- Albert, E., and Podolsky, R. (1935). Can quantum-mechanical description of physical reality be considered complete?
- Asasi, J. *et al*, (2013). Enhanced sensitivity of LIGO gravitational wave detector by using squeezed state of light, *Nature Photonics* **7**, 613.
- Aspect A., Grangier, P., and Roger, G. (1981). Experimental tests of realistic local theories via Bell's theorem *Phys. Rev. Lett.* **47** 460–3
- Aspect A., Grangier, P., and Roger, G. (1982a). Experimental realization of Einstein–Podolsky–Rosen Bohm gedanken experiment: a new violation of Bell's inequalities *Phys. Rev. Lett.* **49** 914
- Aspect A., Grangier, P., and Roger, G. (1982b). Experimental test of Bell's inequality using time varying analyzers *Phys. Rev. Lett.* **49** 1804–7

- Beige, A., Braun, D., Tregenna, B., and Knight, P. L. (2000). Quantum computing using dissipation to remain in a decoherence-free subspace, *Phys. Rev. Lett.* 85, 1762
- Bell, J. S. (1964). On the Einstein–Podolsky–Rosen paradox *Physics* (NY) 1 195–200
- Bennett, C. H., and Brassard, G. (1984). Quantum cryptography: public key distribution and coin tossing *Proc. of the IEEE Int. Conf. on Computer systems and Signal Processing* (Bangalore, India) (Piscataway, NJ : IEEE)
- Bennett, C. H., and Brassard, G. *Sigact News*, vol. 20, no. 4, pp. 78–82, Nov. 1989, <https://doi.org/10.1145/74074.74087>.
- Bennett, C. H., and Brassard, G. (1984). Quantum cryptogra-phy" Public key distribution and coin tossing, *Proc. IEEE International Conf on Computers, Systems and Signal Processing* 175, 8
- Bennett, C. H., Bernstein, H. J., Popescu, S., and Schumacher, B. (1996). Concentrating partial entanglement by local operations, *Phys. Rev. A* 53, 2046
- Bennett, C. H., and Brassard, G. (2014). Quantum cryptography: public key distribution and coin tossing *Theor. Comput. Sci.* 560 7
- Bennett, C. H., Brassard, G., Popescu, S., Schumacher, B., Smolin, J. A., and Woiters, W. K. (1996). "Purification of noisy entanglement and faithful teleportation via noisy channels," *Physical Review Letters*, vol. 76, no. 5, pp. 722–725, 1996.
- Bergquist, J.C., Hulet, R.B., Itano, W.M. and Wineland, D.J. (1986). Observation of Quantum Jumps in a Single Atom, *Phys. Rev. Lett.* 57, 1699
- Bouchée, T., Thurlings, M., De Putter-Smits, L., and Pepin, B. (2023). Investigating teachers' and students' experiences of quantum physics lessons: opportunities and challenges *Res. Sci. Technol. Educ.* 41 777
- Bose, S., Vedral, V., and Knight, P. L. (1999). Purication via entanglement swapping and conserved entanglement, *Phys. Rev. A* 60, 194
- Bose, S., Vedral, V., and Knight, P. L. (1998). Multiparticle gen-eralization of entanglement swapping, *Phys. Rev. A* 57, 822
- Briegel, H.J., Dür, W., Cirac, J.I., and Zoller, P. "Quantum repeaters:: The role of imperfect local operations in quantum communication". *PHYSICAL REVIEW LETTERS* 81, 5932– 5935 (1998).
- Cook, R. J. and Kimble, H. J. (1985). Possibility of direct ob-servation of quantum jumps, *Phys. Rev. Lett.* 54, 1023
- Das, R., Md. S. Rahman and M. Majumdar, "Design of a quantum repeater using quantum circuits and benchmarking its performance on an IBM quantum computer," *Quantum Information Processing*, vol. 20, no. 7, pp. 245, 2021.
- Deutsch, D., Ekert, A., Jozsa, R., Macchiavello, C., Popescu, S., *et al.*, "Quantum privacy amplification and the security of quantum cryptography over noisy channels," *Physical Review Letters*, vol. 77, no. 13, pp. 2818–2821, 1996.
- Deutsch, D. (1985). Quantum theory, the Church-Turing prin-ciple and the universal quantum computer, *Proc. Royal Soc. (London)* 400, 97
- Duarte, F. J. (2022), *Fundamentals of Quantum Entanglement* (Second Edition), IOP Publishing, Bristol, UK,
- Eberly, J. H., Narozhny, N.B., and Sanchez-Mondragon, J.J. (1980). Periodic spontaneous collapse and revival in a simple quantum model, *Phys. Rev. Lett.* 44, 1323
- Einstein, A., Podolsky, B., and Rosen, N. (1935). Can quantum-mechanical description of physical reality be considered complete?, *Phys. Rev.* 47, 777
- Ekert, A. K., and Knight, P. L. (1995). Entangled quantum sys-tems and the Schmidt decomposition, *Am. J. Phys.* 63, 415
- Ekert, A. K. (1991). Quantum cryptography based on Bell's theorem, *Phys. Rev. Lett* 67, 661
- Feynman, R. P. (1982). Simulating physics with computers, *Int. J. Theoretical Phys.* 21, 467
- Filgueiras, J. G., Sarthour, R. S., Souza, A. M., Oliveira, I. S., Serra, R. M., and C'eleri, L. C. Wave-particle duality in an environment with arbitrary white noise, arXiv:1208.0802v2 [quant-ph] 1 Sep 2012
- Fitzgerald, B. W., Emonts, P., and Jordi T. (2024). A Christmas story about quantum teleportation, *Phys. Educ PublishedbyIOPPublishingLtd*, (23pp)
- Gea-Banacloche, J. (1990). Collapse and revival of the state vec-tor in the Jaynes-Cummings model: An

- example of state preparation by a quantum apparatus, *Phys. Rev. Lett.* **65**, 3385
- Gerry, C. C., and Knight, P. L. *Introductory quantum optics* (Cambridge, 2005).
- Ghirardi, G. C., Rimini, A., and Weber, T. (1987). Disentanglement of quantum functions: answer to comment on 'Unified dynamics for microscopic and macroscopic systems' *Phys. Rev. D* **36** 3287
- Gisin, N., et al. (2002). "Quantum cryptography." *Reviews of Modern Physics*.
- Gisin, N., and Thew, R. "Quantum communication," *Nature Photonics*, vol. 1, no. 3, pp. 165–171, 2007.
- Glauber, R. J. (1963). Photon Correlations, *Phys. Rev. Lett.* **10**, 84
- Greenberger, D. M., Horne, M. A., Shimony, A., and Zeilinger, A. (1990). Bell's theorem without inequalities *Am. J. Phys.* **58** 1131–43
- Greinert, F., Müller, R., Bitzenbauer, P., Ubben, M. S., and Weber, K.-A. (2023). Future quantum workforce: competences, requirements and forecasts *Phys. Rev. Phys. Educ. Res.* **19** 010137
- Guang-Liang Li, On Quantum Superposition, July 11, 2023. <https://doi.org/10.32388/N5HMJV>
- Herbst, T., Scheidl, T., Fink, M., Handsteiner, J., Wittmann, B., et al., "Teleportation of entanglement over 143 km," *Proceedings of The National Academy of Sciences*, vol. 112, no. 46, pp. 14202–14205, 2015.
- Hu, X. M., Huang C. X., Sheng, Y. B., Zhou, L., Liu, B. H., et al., "Long-distance entanglement purification for quantum communication," *Physical Review Letters*, vol. 126, no. 1, pp. 010503, 2021. ID quantique, www.idquantique.com.
- Introne, J., Gokce, Y. I., Iandoli, L., DeCook, J., and Elzeini S 2018 How people weave online information into pseudoknowledge *Soc. Media Soc.* **4** 205630511878563
- Jaynes, E. T., and Cummings, F. W. (1963). Comparison of quantum and semiclassical radiation theories with application to the beam maser. *Proc. IEEE.* **51**, 89
- Knight, P. L., and Allen, L. *Concepts of quantum optics* (Pergamon, 1983).
- Knight, P. L., Loudon, R., and Pegg, D. T. (1986). Quantum jumps and atomic cryptograms, *Nature* **323**, 608
- Krastanov, S., Albert, V. V., and Jiang, L. "Optimized entanglement purification," *Quantum*, vol. 3, pp. 123, 2019.
- Krauss, L. M. (1995). *The Physics of Star Trek* (Basic Books)
- Krijtenburg-Lewerissa K., Pol, H. J., Brinkman, A., and Van Joolingen, W. R. (2017). Insights into teaching quantum mechanics in secondary and lower undergraduate education *Phys. Rev. Phys. Educ. Res.* **13** 010109
- Ladd, T. D., Jelezko, F., Laflamme, R., Nakamura, Y., Monroe, C., and O'Brien, J. L. "Quantum computers". *NATURE* **464**, 45–53 (2010).
- Lee, J., Kim, M. S., Park, Y. J., and Lee, S. (2000). Partial tele-transportation of entanglement in a noisy environment, *J. Mod. Opt.* **47**, 2151
- Loudon, R., and Knight, P. L. (1987). Squeezed light, *J. Mod. Opt.* **34**, 709
- Lv, D. et al., Reconstruction of the Jaynes-Cummings eigenstate of ionic motion in a harmonic trap, *Phys. Rev. A* **95**, 043813 (2017).
- Manik, B., Sankar, B., Nirman, G., Tamal, G., Amit, M., Ashutosh, R., and Arup, R. Two-Qubit Pure Entanglement as Optimal Social Welfare Resource in Bayesian Game <https://quantum-journal.org/papers/q-2019-09-09-185/>
- Meekhof, D. M., Monroe, C., King, B. E., Itano, W. M., and Wineland, D. J. (1996). Generation of Nonclassical Motional States of a Trapped Atom, *Phys. Rev. Lett.* **76**, 1796
- Meyer, J. C., Pollock, S. J., Wilcox, B. R., and Passante, G. (2023). How media hype affects our physics teaching: a case study on quantum computing *Phys. Teach.* **61** 339
- Mugambi, K. A., and Okeng'o, G. O. (2023). Design and Implementation of Quantum Repeaters: Insights on Quantum Entanglement Purification, *Journal of Quantum Computing*, 5, 26–40.
- Muralidharan, S., Li, L., Kim, J., Lütkenhaus, N., Lukin, M. D. et al., "Optimal architectures for long distance quantum communication," *Scientific Reports*, vol. 6, no. 1, pp. 20463, 2016.

- Nielsen M A and Chuang I L 2010 *Quantum computation and quantum information* (Cambridge: Cambridge University Press)
- Paspalakis, E., and Knight, P. L. (1998). Phase control of spon-taneous emission, *Phys. Rev. Lett.* 81, 293 4
- Phoenix, S. J. D., and Knight, P. L. (1988). Fluctuations and entropy in models of quantum optical resonance, *Ann Phys.* 186, 381
- Plenio, M. B., and Knight, P. L. (1998).The quantum-jump ap-proach to dissipative dynamics in quantum optics, *Rev. Mod. Phys.* 70, 101
- Rabbie, J., Chakraborty, K., Avis, G., and Wehner, S. “Designing quantum networks using preexisting infrastructure,” *npj Quantum Information*, vol. 8, no. 1, pp. 5, 2022.
- Rempe, G., Walther, H., and Klein, N. (1987). Observation of quantum collapse and revival in a one-atom maser". *Phys. Rev. Lett.* 58, 353
- Ruihong, Q., and Ying, M. “Research progress of quantum repeaters,” *Journal of Physics: Conference Series*, vol. 1237, pp. 5, 2019.
- Scarani, V., Bechmann-Pasquinucci, H., Cerf, N. J., Dušek, M., Lütkenhaus, N., and M. Peev, *Rev. Mod. Phys.*, vol. 81, no. 3, pp. 1301–1350, Sept. 2009, doi: <https://doi.org/10.1103/RevModPhys.81.1301>.
- Shor, P. W. (1997). Polynomial-time algorithms for prime fac-torization and discrete logarithms on a quantum com-puter, *SIAM J. Comput.*, 26, 1484
- Shor, P. W. (1995). Scheme for reducing decoherence in quan-tum computer memory, *Phys. Rev. A* 52, R2493
- Shore, B. W., and Knight, P. L. (1993). The Jaynes-Cummings model, *J. Mod. Opt.* 40, 1195
- Shor, P. W. in *35th Annual Symposium on Foundations of Computer Science*, pp. 124–134, Nov. 1994, doi: <https://doi.org/10.1109/SFCS.1994.365700>.
- Shor, P. W. *SIAM Journal of Computing*, vol. 26, no. 5, pp. 1484–1509, Oct. 1997, doi: <https://doi.org/10.1137/S0097539795293172> .
- Singh, C., Levy, A., and Levy, J. (2022). Preparing precollege students for the second quantum revolution with core concepts in quantum information science *Phys. Teach.* 60 639
- Sutrini, C., Malgieri, M., Zuccarini, G., and Macchiavello, C. (2023). A teacher professional development course on quantum technologies: discussion of results *J. Phys.: Conf. Ser.* 2490 012006
- Sridhar G T., Ashwini P., and Tabassum, N. (2023). A review on quantum communication and computing 2023 *2nd Int. Conf. on Applied Artificial Intelligence and Computing (ICAAIC)* pp 1592–6
- Steane, A. M. (1999). Ecient fault-tolerant quantum comput-ing, *Nature* 399, 124
- Sudarshan, E. C. G. (1963). Equivalence of Semiclassical and Quantum Mechanical Descriptions of Statistical Light Beams, *Phys. Rev. Lett.* 10, 277
- Travis, S. H. (2014). [A Overview of Quantum Teleportation: A Technology for Potential Intelligence-gathering Applications](https://www.researchgate.net/publication/266111111_A_Overview_of_Quantum_Teleportation:_A_Technology_for_Potential_Intelligence-gathering_Applications). Researchgate
- Tsai, C.-W.; Yang, C.-W.; Lin, J.; Chang, Y.-C.; Chang, R.-S. Quantum Key Distribution Networks: Challenges and Future Research Issues in Security. *Appl. Sci.* 2021, 11, 3767. <https://doi.org/10.3390/app1109376>
- Ulf, L. (2010). *Essential Quantum Optics: From Quantum Measurements to Black Holes*. Cambridge University Press, New York, United States of America.www.cambridge.org
- Vedral, V., Plenio, M. B., Rippin, M. A., and Knight, P. L. (1997). Quantifying entanglement, *Phys. Rev. Lett.* 78 2275
- Vedral, V., and Plenio, M. B. (1998). Entanglement measures and purification procedures, *Phys. Rev. A* 57, 1619
- Vicente M., Jesus M., and Momtchil P. (2016). Introduction to Quantum Key Distribution. Center for Computational Simulation, Universidad Polit’ecnica de Madrid, 28660 Boadilla del Monte, Madrid, Spain
- Wei S-H *et al* 2022 Towards real-world quantum networks: a review *Laser Photonics Rev.* 16 2100219
- Wehner S, Elkouss D and Hanson R 2018 Quantum internet: a vision for the road ahead *Science* 362 eaam9288
- Yihe-Xue, Basic Theory of Quantum Entanglement and the Possibility of Passing on Information Faster than the Speed of Light, 3rd International Forum on Geoscience

and Geodesy, iop publication, IOP Conf. Series: Earth and Environmental Science, 658, (2021) 012001

Yin J *et al* 2017 Satellite-based entanglement distribution over 1200 kilometers *Science* **356** 1140

Yuen, H., and Shapiro, J. H. (1978). Optical communications with two-photon coherent states - part I: Quantum state propagation and quantum-noise reduction, *IEEE Trans Inform Theory* IT 24, 657

Zapatero, V., van Leent, T., Arnon-Friedman, R., Liu, W-Z., Zhang, Q., Weinfurter, H., and Curty, M. (2023). Advances in device-independent quantum key distribution *npj Quantum Inf.* **9** 1

Zukowski, M., Zeilinger, A., Horne, M. A., and Ekert, A. K. (1993). 'Event-ready-detectors' Bell experiment via entanglement swapping, *Phys. Rev. Lett.* 71, 4287